

GENERALIZATION OF n -ARY NAMBU ALGEBRAS AND BEYOND

H. ATAGUEMA, A. MAKHLOUF AND S. SILVESTROV

ABSTRACT. The aim of this paper is to introduce n -ary Hom-algebra structures generalizing the n -ary algebras of Lie type enclosing n -ary Nambu algebras, n -ary Nambu-Lie algebras, n -ary Lie algebras, and n -ary algebras of associative type enclosing n -ary totally associative and n -ary partially associative algebras. Also, we provide a way to construct examples starting from an n -ary algebra and an n -ary algebras endomorphism. Several examples could be derived using this process.

INTRODUCTION

In this paper, we introduce generalizations of n -ary algebras of Lie type and associative type by twisting the identities using linear maps. The n -ary algebraic structures and in particular ternary algebraic structures appeared more or less naturally in various domains of theoretical and mathematical physics and data processing. Indeed, theoretical physics progress of quantum mechanics and the discovery of the Nambu mechanics (1973) (see [68]), as well as a work of S. Okubo (see [69]) on Yang-Baxter equation gave impulse to a significant development on n -ary algebras. The n -ary operations appeared first through cubic matrices which were introduced in the nineteenth century by Cayley. The cubic matrices were considered again and generalized by Kapranov, Gelfand, Zelevinskii in 1994 (see [43]) and Sokolov in 1972 (see [75]). Another recent motivation to study n -ary operation comes from string theory and M-branes involving naturally an algebra with ternary operation called Bagger-Lambert algebra [5]. Hundred of papers are dedicated to Bagger-Lambert algebra. For other physical applications (see [46, 47, 48, 49]).

The first conceptual generalization of binary algebras was the ternary algebras introduced by Jacobson (see [42]). In connection with problems from Jordan theory and quantum mechanics, he defined the Lie triple systems. A Lie triple system consists of a space of linear operators on vector space V that is closed under the ternary bracket $[x, y, z]_T = [[x, y], z]$, where $[x, y] = xy - yx$. Equivalently, the Lie triple system may be viewed as a subspace of the Lie algebra closed relative to the ternary product. A Lie triple system arose also in the study of symmetric spaces (see [60]). More generally, we distinguish two kinds of generalizations of binary Lie algebras. Firstly, n -ary Lie algebras in which the Jacobi identity is generalized by considering a cyclic summation over $\mathcal{S}_{2n-1}$ instead of $\mathcal{S}_3$, (see [34] [66]) and secondly n -ary Nambu algebras in which the fundamental identity generalizes the fact that the adjoint maps are derivations. The fundamental identity appeared first in Nambu mechanics [68], the abstract definition of n -ary Nambu algebras or n -ary Nambu-Lie algebras (when the bracket is skew symmetric) was given by Fillipov in 1985 (see [24]). See also ([76, 77]) for the algebraic formulation of the Nambu mechanics. The Leibniz n -ary algebras were introduced and studied in [11]. While the basic concepts of ternary Hopf algebra were introduced in [74] and [8], see also [31].

On the other hand, n -ary algebras of associative type were studied by Lister, Loos, Myung and Carlsson (see [9, 57, 59, 67]). The n -ary operations of associative type lead to two principal classes of "associative" n -ary algebras, totally associative n -ary algebras and partially associative n -ary algebras. Also they admit some variants. Typical examples of totally associative n -ary algebras for $n = 3$ are given by subspaces of

Date: November 2008.

2000 Mathematics Subject Classification. 17A30, 17A40, 17A42, 17D99.

Key words and phrases. n -ary totally Hom-Associative algebra, n -ary partially Hom-Associative algebra, n -ary Hom-Lie algebra, n -ary Hom-Nambu algebra, n -ary Hom-Nambu-Lie algebra.

This work was partially supported by the Crafoord Foundation, The Swedish Foundation for International Cooperation in Research and Higher Education (STINT), The Swedish research links program of SIDA foundation and the Swedish Research Council, The Royal Swedish Academy of Sciences, The Royal Physiographic Society in Lund, The European network Liegrits, Mulhouse University and Lund University.

an associative algebra that are closed relative to the ternary product $(xyz) \mapsto xyz$. This natural ternary product is related to the important ternary operation introduced by Hestenes (see [38]) defined on the linear space of rectangular matrices $A, B, C \in \mathcal{M}_{m,n}$ with complex entries by AB^*C where B^* is the conjugate transpose matrix of B . This operation is strictly speaking not a ternary algebra product on $\mathcal{M}_{m,n}$ as it is linear on the first and the third arguments but conjugate-linear on the second argument. It satisfies identities, sometimes referred to as identities of total associativity of second kind, only slightly different from the identities of totally associative algebras. The totally associative ternary algebras are also sometimes called associative triple systems. The cohomology of totally associative n -ary algebras was studied by Carlsson through the embedding (see [10]). In [3], the first and second authors extended the 1-parameter formal deformation theory to ternary algebras of associative type and in [4] they discussed their cohomologies adapted to deformations, see also [39, 30, 73].

The Hom-algebras structures arose first in quasi-deformation of Lie algebras of vector fields. These quasi-deformations lead to quasi-Lie algebras, a generalized Lie algebra structure in which the skew-symmetry and Jacobi conditions are twisted. The first examples were concerned with q -deformations of the Witt and Virasoro algebras (see [2, 12, 13, 14, 15, 16, 18, 44, 58, 40]). Motivated by these and the new examples arising as application of the general quasi-deformation construction of [36, 54, 55] on the one hand, and the desire to be able to treat within the same framework such well-known generalizations of Lie algebras as the color and super Lie algebras on the other hand, quasi-Lie algebras and subclasses of quasi-Hom-Lie algebras and Hom-Lie algebras were introduced in [36, 54, 55, 56]. In the subclass of Hom-Lie algebras skew-symmetry is untwisted, whereas the Jacobi identity is twisted by a single linear map and contains three terms as in Lie algebras, reducing to ordinary Lie algebras when the twisting linear map is the identity map.

The notion of Hom-associative algebras generalizing associative algebras to a situation where associativity law is twisted by a linear map was introduced by the second and the third authors in [61], where it was shown in particular that the commutator product, defined using the multiplication in a Hom-associative algebra, leads naturally to a Hom-Lie algebra. We introduced also Hom-Lie-admissible algebras and more general G -Hom-associative algebras with subclasses of Hom-Vinberg and pre-Hom-Lie algebras, generalizing to the twisted situation Lie-admissible algebras, G -associative algebras, Vinberg and pre-Lie algebras respectively, and shown that for these classes of algebras the operation of taking commutator leads to Hom-Lie algebras as well. The enveloping algebras of Hom-Lie algebras were discussed in [79]. It is important challenging problem to develop further the proper notion and theory of universal enveloping algebras for general Hom-algebras and for general quasi-Hom-Lie and Quasi-Lie algebras. The fundamentals of the formal deformation theory and associated cohomology structures for Hom-Lie algebras have been considered recently by the second and the third authors in [63]. Simultaneously, D. Yau has developed elements of homology for Hom-Lie algebras in [80]. In [62] and [64], we developed the theory of Hom-coalgebras and related structures. We introduced Hom-coalgebra structure, leading to the notions of Hom-bialgebra and Hom-Hopf algebra, proved some fundamental properties and provided examples. Also, we defined the concept of Hom-Lie admissible Hom-coalgebra generalizing the admissible coalgebra introduced in [29], and provide their classification based on subgroups of the symmetric group. This paper is organized as follows. Section 1 is dedicated to a generalization of the n -ary algebras of Lie type to Hom structures. We introduce the notions of n -ary Hom-Nambu algebra, n -ary Hom-Nambu-Lie algebra and n -ary Hom-Lie algebra. Also, we provide a process to construct them starting from n -ary Nambu algebra (resp. n -ary Nambu-Lie algebra or n -ary Lie algebra) and an algebra endomorphism of this n -ary algebra. In Section 2, we introduce the n -ary Hom-algebras of associative type, and provide a procedure for construction of such n -ary Hom-algebras from n -ary algebras of the same associative type and their n -ary algebra endomorphisms. We consider the totally Hom-associative n -ary algebras and partially Hom-associative n -ary algebras. Their variants, a weak totally associative n -ary algebras and the alternate partially associative n -ary algebras, could be also considered in the same scheme. We provide several examples, using this procedure. In Section 3, we present some results concerning the tensor product of n -ary Hom-algebras.

1. THE n -ARY HOM-ALGEBRAS OF LIE TYPE

Throughout this paper, we will for simplicity of exposition assume that $\mathbb{K}$ is an algebraically closed field of characteristic zero, even though for most of the general definitions and results in the paper this assumption is not essential. Let V be a $\mathbb{K}$ -vector space.

1.1. Definitions. In this section, we introduce the definition of n -ary Hom-Nambu algebras, n -ary Hom-Nambu-Lie algebras and n -ary Hom-Lie algebras, generalizing the usual n -ary Nambu algebras, n -ary Nambu-Lie algebras (called also Filippov n -ary algebras) and n -ary Lie algebras.

Definition 1.1. An n -ary Hom-Nambu algebra is a triple $(V, [\dots], \alpha)$, consisting of a vector space V , an n -linear map $[\dots] : V^{\times n} \rightarrow V$ and a family $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ of linear maps $\alpha_i : V \rightarrow V$, $i = 1, \dots, n-1$ satisfying

$$(1.1) \quad [\alpha_1(x_1), \dots, \alpha_{n-1}(x_{n-1}), [x_n, \dots, x_{2n-1}]] = \sum_{i=n}^{2n-1} [\alpha_1(x_n), \dots, \alpha_{i-n}(x_{i-1}), [x_1, \dots, x_{n-1}, x_i], \alpha_{i-n+1}(x_{i+1}), \dots, \alpha_{n-1}(x_{2n-1})]$$

for all $(x_1, \dots, x_{2n-1}) \in V^{2n-1}$.

We call the condition (1.1) the n -ary Hom-Nambu identity.

In particular, the ternary Hom-Nambu algebras are defined by the following Hom-Nambu identity

$$(1.2) \quad [\alpha_1(x_1), \alpha_2(x_2), [x_3, x_4, x_5]] = [[x_1, x_2, x_3], \alpha_1(x_4), \alpha_2(x_5)] + [\alpha_1(x_3), [x_1, x_2, x_4], \alpha_2(x_5)] + [\alpha_1(x_3), \alpha_2(x_4), [x_1, x_2, x_5]].$$

Remark 1.2. Let $(V, [\dots], \alpha)$ be an n -ary Hom-Nambu algebra where $\alpha = (\alpha_i)_{i=1, \dots, n-1}$. For any $x = (x_1, \dots, x_{n-1}) \in V^{n-1}$, $\alpha(x) = (\alpha_1(x_1), \dots, \alpha_{n-1}(x_{n-1})) \in V^{n-1}$ and $y \in V$, let L_x be a linear map on V , defined by

$$L_x(y) = [x_1, \dots, x_{n-1}, y].$$

Then the Hom-Nambu identity may be written as

$$L_{\alpha(x)}([x_n, \dots, x_{2n-1}]) = \sum_{i=n}^{2n-1} [\alpha_1(x_n), \dots, \alpha_{i-n}(x_{i-1}), L_x(x_i), \alpha_{i-n+1}(x_{i+1}), \dots, \alpha_{n-1}(x_{2n-1})].$$

Remark 1.3. When the maps $(\alpha_i)_{i=1, \dots, n-1}$ are all identity maps, one recovers the classical n -ary Nambu algebras. In the special case of classical n -ary Nambu algebra, it's known also as the fundamental identity or Filippov identity.

Definition 1.4. An n -ary Hom-Nambu algebra $(V, [\dots], \alpha)$ where $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ is called n -ary Hom-Nambu-Lie algebra if the bracket is skew-symmetric that is

$$(1.3) \quad [x_{\sigma(1)}, \dots, x_{\sigma(n)}] = \text{Sgn}(\sigma)[x_1, \dots, x_n], \quad \forall \sigma \in \mathcal{S}_n \text{ and } \forall x_1, \dots, x_n \in V$$

where $\mathcal{S}_n$ stands for the permutation group on n elements.

Definition 1.5. An n -ary Hom-Lie algebra is a triple $(V, [\dots], \alpha)$, consisting of a vector space V , a skew-symmetric map $[\dots] : V^{\times n} \rightarrow V$ and a family $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ of linear maps $\alpha_i : V \rightarrow V$, $i = 1, \dots, n-1$ satisfying n -ary Hom-Jacobi identity

$$(1.4) \quad \sum_{\sigma \in \mathcal{S}_{2n-1}} \text{Sgn}(\sigma) [\alpha_1(x_{\sigma(1)}), \dots, \alpha_{n-1}(x_{\sigma(n-1)}), [x_{\sigma(n)}, \dots, x_{\sigma(2n-1)}]] = 0$$

for all $(x_1, \dots, x_{2n-1}) \in V^{2n-1}$.

In particular, the ternary Hom-Lie algebras are defined by the following ternary Hom-Jacobi identity

$$\sum_{\sigma \in \mathcal{S}_5} \text{Sgn}(\sigma) [\alpha_1(x_{\sigma(1)}), \alpha_2(x_{\sigma(2)}), [x_{\sigma(3)}, x_{\sigma(4)}, x_{\sigma(5)}]] = 0.$$

The morphisms of n -ary Hom-algebras of Lie type are defined as follows.

Definition 1.6. Let $(V, [\cdot, \dots, \cdot], \alpha)$ and $(V', [\cdot, \dots, \cdot]', \alpha')$ be two n -ary Hom-Nambu algebras (resp. n -ary Hom-Nambu-Lie algebras, n -ary Hom-Lie algebras) where $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ and $\alpha' = (\alpha'_i)_{i=1, \dots, n-1}$. A linear map $\rho : V \rightarrow V'$ is an n -ary Hom-Nambu algebras morphism (resp. n -ary Hom-Nambu-Lie algebras morphism, n -ary Hom-Lie algebras morphism) if it satisfies

$$\begin{aligned} \rho([x_1, \dots, x_n]) &= [\rho(x_1), \dots, \rho(x_n)]' \\ \rho \circ \alpha_i &= \alpha'_i \circ \rho \quad \forall i = 1, \dots, n-1. \end{aligned}$$

1.2. Construction of n -ary Hom-algebras of Lie type. The following theorem provides a way to construct an n -ary Hom-Nambu algebra (resp. n -ary Hom-Nambu-Lie algebra, n -ary Hom-Lie algebra) starting from an n -ary Nambu algebra (resp. n -ary Nambu-Lie algebra, n -ary Lie algebra) and an n -ary algebras endomorphism.

Theorem 1.7. Let (V, m) be an n -ary Nambu algebra (resp. n -ary Nambu-Lie algebra, n -ary Lie algebra) and let $\rho : V \rightarrow V$ be an n -ary Nambu (resp. n -ary Nambu-Lie, n -ary Lie) algebras endomorphism. We set $m_\rho = \rho \circ m$ and $\tilde{\rho} = (\rho, \dots, \rho)$. Then $(V, m_\rho, \tilde{\rho})$ is an n -ary Hom-Nambu (resp. n -ary Hom-Nambu-Lie, n -ary Hom-Lie) algebra.

Moreover, suppose that (V', m') is another n -ary Nambu algebra (resp. n -ary Nambu-Lie algebra, n -ary Lie algebra) and $\rho' : V' \rightarrow V'$ is a n -ary Nambu (resp. n -ary Nambu-Lie, n -ary Lie) algebra endomorphism. If $f : V \rightarrow V'$ is a n -ary Nambu algebra morphism (resp. n -ary Nambu-Lie algebra morphism, n -ary Lie algebra morphism) that satisfies $f \circ \rho = \rho' \circ f$ then

$$f : (V, m_\rho, \tilde{\rho}) \longrightarrow (V', m'_{\rho'}, \tilde{\rho}')$$

is an n -ary Hom-Nambu algebras morphism (resp. n -ary Hom-Nambu-Lie algebras morphism, n -ary Hom-Lie algebras morphism).

Proof. We show that $(V, m_\rho, \tilde{\rho})$ satisfies the Hom-Nambu identity. Indeed

$$\begin{aligned} m_\rho(\rho(x_1), \dots, \rho(x_{n-1}), m_\rho(x_n, \dots, x_{2n-1})) &= \rho \circ m(\rho(x_1), \dots, \rho(x_{n-1}), \rho \circ m(x_n, \dots, x_{2n-1})) \\ &= \rho^2 \circ m(x_1, \dots, x_{n-1}, m(x_n, \dots, x_{2n-1})). \end{aligned}$$

On the other hand,

$$\begin{aligned} &\sum_{i=1}^n m_\rho(\rho(x_n), \dots, \rho(x_{i-1}), m_\rho(x_1, \dots, x_{n-1}, x_i), \rho(x_{i+1}), \dots, \rho(x_{2n-1})) \\ &= \sum_{i=1}^n \rho \circ m(\rho(x_n), \dots, \rho(x_{i-1}), \rho \circ m(x_1, \dots, x_{n-1}, x_i), \rho(x_{i+1}), \dots, \rho(x_{2n-1})) \\ &= \rho^2 \left(\sum_{i=1}^n m(x_n, \dots, x_{i-1}, m(x_1, \dots, x_{n-1}, x_i), x_{i+1}, \dots, x_{2n-1}) \right). \end{aligned}$$

Therefore the Nambu identity implies the Hom-Nambu identity for this n -ary algebra. The skew-symmetry and the Hom-Jacobi identity are proved similarly. The second assertion follows from

$$f \circ m_\rho = f \circ \rho \circ m = \rho' \circ f \circ m = \rho' \circ m' \circ f = m'_{\rho'} \circ f.$$

□

Example 1.8. An algebra V consisting of polynomials or possibly of other differentiable functions in 3 variables x_1, x_2, x_3 , equipped with well-defined bracket multiplication given by the functional jacobian $J(f) = (\frac{\partial f_i}{\partial x_j})_{1 \leq i, j \leq 3}$:

$$(1.5) \quad [f_1, f_2, f_3] = \det \begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \frac{\partial f_1}{\partial x_3} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \frac{\partial f_2}{\partial x_3} \\ \frac{\partial f_3}{\partial x_1} & \frac{\partial f_3}{\partial x_2} & \frac{\partial f_3}{\partial x_3} \end{pmatrix},$$

is a ternary Nambu-Lie algebra. By considering a ternary Nambu-Lie algebra endomorphism of such algebra, we construct a ternary Hom-Nambu-Lie algebra. Let $\gamma(x_1, x_2, x_3)$ be a polynomial or more

general differentiable transformation of three variables mapping elements of V to elements of V and such that $\det J(\gamma) = 1$. Let $\rho_\gamma : V \mapsto V$ be the composition transformation defined by $f \mapsto f \circ \gamma$ for any $f \in V$. By the general chain rule for composition of transformations of several variables,

$$\begin{aligned} J(\rho_\gamma(f)) &= J(f \circ \gamma) = (J(f) \circ \gamma)J(\gamma) = \rho_\gamma(J(f))J(\gamma), \\ \det J(\rho_\gamma(f)) &= \det(J(f) \circ \gamma) \det J(\gamma) = \det \rho_\gamma(J(f)) \det J(\gamma). \end{aligned}$$

Hence, for any transformation γ with $\det J(\gamma) = 1$, the composition transformation ρ_γ defines an endomorphism of the ternary Nambu-Lie algebra with ternary product (1.5). Therefore, by Theorem 1.7, for any such transformation γ , the triple $(V, [\cdot, \cdot, \cdot]_\gamma = \rho_\gamma \circ [\cdot, \cdot, \cdot], \widetilde{\rho}_\gamma)$ is a ternary Hom-Nambu-Lie algebra.

2. THE n -ARY HOM-ALGEBRAS OF ASSOCIATIVE TYPE

In this Section, we introduce the n -ary totally Hom-associative algebras and n -ary partially Hom-associative algebras generalizing the classical n -ary totally associative algebras and n -ary partially associative algebras. We also provide a process to construct them starting from an n -ary totally associative algebra or n -ary partially associative algebra and an algebra endomorphism of such n -ary algebra.

2.1. Definitions. We introduce the definitions of n -ary totally Hom-associative algebras and n -ary partially Hom-associative algebras.

Definition 2.1. An n -ary totally Hom-associative algebra is a triple (V, m, α) , consisting of a vector space V , a n -linear map $m : V^{\times n} \rightarrow V$ and a family $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ of linear maps $\alpha_i : V \rightarrow V$, $i = 1, \dots, n-1$ satisfying

$$\begin{aligned} (2.1) \quad & m(\alpha_1(x_1), \dots, \alpha_{n-1}(x_{n-1}), m(x_n, \dots, x_{2n-1})) = \\ & \quad \quad \quad \vdots \\ & m(\alpha_1(x_1), \dots, \alpha_i(x_i), m(x_{i+1}, \dots, x_{i+n}), \alpha_{i+1}(x_{i+1+n}), \dots, \alpha_{n-1}(x_{2n-1})) = \\ & \quad \quad \quad \vdots \\ & m(m(x_1, \dots, x_n), \alpha_1(x_{n+1}), \dots, \alpha_{n-1}(x_{2n-1})). \end{aligned}$$

Definition 2.2. An n -ary partially Hom-associative algebra is a triple (V, m, α) , consisting of a vector space V , a n -linear map $m : V^{\times n} \rightarrow V$ and a family $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ of linear maps $\alpha_i : V \rightarrow V$, $i = 1, \dots, n-1$ satisfying

$$(2.2) \quad \sum_{i=0}^{n-1} m(\alpha_1(x_1), \dots, \alpha_i(x_i), m(x_{i+1}, \dots, x_{i+n}), \alpha_{i+1}(x_{i+1+n}), \dots, \alpha_{n-1}(x_{2n-1})) = 0.$$

Remark 2.3. When the maps $(\alpha_i)_{i=1, \dots, n-1}$ are all identity maps then one recovers the classical n -ary totally or partially associative algebras.

Remark 2.4. These generalizations to n -ary Hom-algebra structures could be extend obviously to the variants of n -ary algebras of associative type. An n -ary weak totally Hom-associative algebras, is given by the identity

$$m(\alpha_1(x_1), \dots, \alpha_{n-1}(x_{n-1}), m(x_n, \dots, x_{2n-1})) = m(m(x_1, \dots, x_n), \alpha_1(x_{n+1}), \dots, \alpha_{n-1}(x_{2n-1})).$$

Also, the n -ary alternate partially Hom-associative algebras, where some signs in the identity (2.2) are moved to minus, could be defined in the same way.

The morphisms of n -ary Hom-algebras of associative type are defined as follows.

Definition 2.5. Let (V, m, α) and (V', m', α') be two n -ary totally Hom-associative algebras (resp. n -ary partially Hom-associative algebras) where $\alpha = (\alpha_i)_{i=1, \dots, n-1}$ and $\alpha' = (\alpha'_i)_{i=1, \dots, n-1}$. A linear map $\rho : V \rightarrow V'$ is an n -ary totally Hom-associative algebras morphism (resp. partially Hom-associative algebras morphism) if it satisfies

$$\rho(m(x_1, \dots, x_n)) = m'(\rho(x_1), \dots, \rho(x_n)) \quad \text{and} \quad \rho \circ \alpha_i = \alpha'_i \circ \rho \quad \forall i = 1, \dots, n-1.$$

2.2. Construction of n -ary Hom-algebras of Associative type. The following theorem provide a way to construct an n -ary totally Hom-associative algebra (resp. n -ary partially Hom-associative algebra) starting from an n -ary totally associative (resp. n -ary partially associative algebra) and n -ary algebras endomorphism.

Theorem 2.6. *Let (V, m) be an n -ary totally associative algebra (resp. n -ary partially associative algebra) and let $\rho : V \rightarrow V$ be an n -ary totally Hom-associative (resp. n -ary partially Hom-associative) algebra endomorphism. We set $m_\rho = \rho \circ m$ and $\tilde{\rho} = (\rho, \dots, \rho)$. Then $(V, m_\rho, \tilde{\rho})$ is an n -ary totally Hom-associative algebra (resp. n -ary partially Hom-associative algebra).*

Moreover, suppose that (V', m') is another n -ary totally associative algebra (resp. n -ary partially associative algebra) and $\rho' : V' \rightarrow V'$ is a n -ary totally associative (resp. n -ary partially associative) algebras endomorphism. If $f : V \rightarrow V'$ is a n -ary totally associative algebra morphism (resp. n -ary partially associative algebra morphism) that satisfies $f \circ \rho = \rho' \circ f$, then

$$f : (V, m_\rho, \tilde{\rho}) \longrightarrow (V', m'_{\rho'}, \tilde{\rho}')$$

is an n -ary totally Hom-associative algebras morphism (resp. n -ary partially Hom-associative algebras morphism).

Proof. We show that $(V, m_\rho, \tilde{\rho})$ satisfies the identity (2.1). Indeed, for $i = 1, \dots, n-1$ we have

$$\begin{aligned} m_\rho(\rho(x_1), \dots, \rho(x_i), m_\rho(x_{i+1}, \dots, x_{i+n}), \rho(x_{i+1+n}), \dots, \rho(x_{2n-1})) &= \\ \rho \circ m(\rho(x_1), \dots, \rho(x_i), \rho \circ m(x_{i+1}, \dots, x_{i+n}), \rho(x_{i+1+n}), \dots, \rho(x_{2n-1})) &= \\ \rho^2(m(x_1, \dots, x_i, m(x_{i+1}, \dots, x_{i+n}), x_{i+1+n}, \dots, x_{2n-1})). \end{aligned}$$

Therefore the n -ary total associativity identity implies the n -ary total Hom-associativity identity for this n -ary algebra. The proof for n -ary partially Hom-associative algebras is similar. The second assertion is proved in the same way as the second part of Theorem 1.7. $\square$

2.3. Examples.

Example 2.7. *In dimension 2, there exists a partially associative ternary algebra defined with respect to a basis $\{e_1, e_2\}$ as follows*

$$m(e_1, e_1, e_1) = e_2, \quad m(e_i, e_j, e_k) = 0, \text{ when } 2 \in \{i, j, k\}.$$

Its automorphisms are given with respect to the same basis by the matrices of the form : $\begin{pmatrix} a & 0 \\ b & a^3 \end{pmatrix}$ with $a, b \in \mathbb{K}$ and $a \neq 0$.

We deduce using Theorem 2.6 the 2-dimensional partially Hom-associative ternary algebra $(\mathbb{K}^2, \tilde{m}, \rho)$ given with respect to a basis $\{e_1, e_2\}$, by the following nontrivial product

$$\tilde{m}(e_1, e_1, e_1) = a^3 e_2,$$

and the linear map

$$\begin{aligned} \rho(e_1) &= a e_1 + b e_2 \\ \rho(e_2) &= a^3 e_2 \end{aligned}$$

Example 2.8. *We consider now the 2-dimensional totally associative ternary algebra $(\mathbb{K}^2, m)$ given with respect to a basis e_1, e_2 of $\mathbb{K}^2$, by the following products :*

$$\begin{aligned} m(e_1, e_1, e_1) &= e_1 & m(e_2, e_2, e_1) &= e_1 + e_2 \\ m(e_1, e_1, e_2) &= e_2 & m(e_2, e_2, e_2) &= e_1 + 2e_2 \\ m(e_1, e_2, e_2) &= e_1 + e_2 & m(e_1, e_2, e_1) &= e_2 \\ m(e_2, e_1, e_1) &= e_2 & m(e_2, e_1, e_2) &= e_1 + e_2 \end{aligned}$$

The following matrices correspond to automorphisms of the previous totally associative ternary algebra.

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \quad \begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix} \\ \begin{pmatrix} -\frac{1}{\sqrt{5}} & -\frac{3}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix} \quad \begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix} \quad \begin{pmatrix} \frac{1}{\sqrt{5}} & \frac{3}{\sqrt{5}} \\ -\frac{2}{\sqrt{5}} & -\frac{1}{\sqrt{5}} \end{pmatrix}$$

We deduce for example using Theorem 2.6 the following 2-dimensional totally Hom-associative ternary algebras $(\mathbb{K}^2, \tilde{m}_i, \rho_i)$ for $i = 1, 2$, which are not totally associative. They are given with respect to a basis $\{e_1, e_2\}$, by the following product

$$\begin{aligned} \tilde{m}_1(e_1, e_1, e_1) &= e_1 & \tilde{m}_1(e_2, e_2, e_1) &= 2e_1 - e_2 \\ \tilde{m}_1(e_1, e_1, e_2) &= e_1 - e_2 & \tilde{m}_1(e_2, e_2, e_2) &= 3e_1 - 2e_2 \\ \tilde{m}_1(e_1, e_2, e_2) &= 2e_1 - e_2 & \tilde{m}_1(e_1, e_2, e_1) &= e_1 - e_2 \\ \tilde{m}_1(e_2, e_1, e_1) &= e_1 - e_2 & \tilde{m}_1(e_2, e_1, e_2) &= 2e_1 - e_2 \end{aligned}$$

and the linear map

$$\begin{aligned} \rho_1(e_1) &= e_1 \\ \rho_1(e_2) &= e_1 - e_2 \end{aligned}$$

The second example is given by

$$\begin{aligned} \tilde{m}_2(e_1, e_1, e_1) &= \frac{1}{\sqrt{5}}e_1 - \frac{2}{\sqrt{5}}e_2 & \tilde{m}_2(e_2, e_2, e_1) &= \frac{4}{\sqrt{5}}e_1 - \frac{3}{\sqrt{5}}e_2 \\ \tilde{m}_2(e_1, e_1, e_2) &= \frac{3}{\sqrt{5}}e_1 - \frac{1}{\sqrt{5}}e_2 & \tilde{m}_2(e_2, e_2, e_2) &= \frac{7}{\sqrt{5}}e_1 - \frac{4}{\sqrt{5}}e_2 \\ \tilde{m}_2(e_1, e_2, e_2) &= \frac{4}{\sqrt{5}}e_1 - \frac{3}{\sqrt{5}}e_2 & \tilde{m}_2(e_1, e_2, e_1) &= \frac{3}{\sqrt{5}}e_1 - \frac{1}{\sqrt{5}}e_2 \\ \tilde{m}_2(e_2, e_1, e_1) &= \frac{3}{\sqrt{5}}e_1 - \frac{1}{\sqrt{5}}e_2 & \tilde{m}_2(e_2, e_1, e_2) &= \frac{4}{\sqrt{5}}e_1 - \frac{3}{\sqrt{5}}e_2 \end{aligned}$$

and the linear map

$$\begin{aligned} \rho_2(e_1) &= \frac{1}{\sqrt{5}}e_1 - \frac{2}{\sqrt{5}}e_2 \\ \rho_2(e_2) &= \frac{3}{\sqrt{5}}e_1 - \frac{1}{\sqrt{5}}e_2 \end{aligned}$$

3. TENSOR PRODUCTS

In this section we define the tensor product of two n -ary Hom-algebras and prove some results involving n -ary Hom-algebras of Lie type and associative type.

Let $\mathcal{A} = (V, m, \alpha)$, where $\alpha = (\alpha_i)_{i=1, \dots, n-1}$, and $\mathcal{A}' = (V', m', \alpha')$, $\alpha' = (\alpha'_i)_{i=1, \dots, n-1}$, be two n -ary Hom-algebras of given type. The tensor product $\mathcal{A} \otimes \mathcal{A}'$ is an n -ary algebra defined by the triple $(V \otimes V', m \otimes m', \alpha \otimes \alpha')$ where $\alpha \otimes \alpha' = (\alpha_i \otimes \alpha'_i)_{i=1, \dots, n-1}$ with

$$\begin{aligned} m \otimes m'(x_1 \otimes x'_1, \dots, x_n \otimes x'_n) &= m(x_1, \dots, x_n) \otimes m'(x'_1, \dots, x'_n), \\ \alpha_i \otimes \alpha'_i(x \otimes x') &= \alpha_i(x) \otimes \alpha'_i(x'), \end{aligned}$$

where $x, x_1, \dots, x_n \in V$ and $x', x'_1, \dots, x'_n \in V'$.

Recall that an n -ary multiplication m is symmetric if

$$(3.1) \quad m(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = m(x_1, \dots, x_n), \quad \forall \sigma \in \mathcal{S}_n \text{ and } \forall x_1, \dots, x_n \in V$$

and m' is skew-symmetric if

$$(3.2) \quad m'(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = \text{Sgn}(\sigma)m'(x_1, \dots, x_n), \quad \forall \sigma \in \mathcal{S}_n \text{ and } \forall x_1, \dots, x_n \in V'$$

where $\mathcal{S}_n$ stands for the permutation group on n elements.

We have the following obvious lemma.

Lemma 3.1. *Let $\mathcal{A} = (V, m, \alpha)$ and $\mathcal{A}' = (V', m', \alpha')$ be two n -ary Hom-algebras of given type. If m is symmetric and m' is skew-symmetric then $m \otimes m'$ is skew-symmetric.*

Proposition 3.2. *Let $\mathcal{A}$ be an n -ary totally Hom-associative algebra and $\mathcal{A}'$ be an n -ary partially Hom-associative algebra, then $\mathcal{A} \otimes \mathcal{A}'$ is an n -ary partially Hom-associative algebra.*

Proof. Using the n -ary totally Hom-associativity of $\mathcal{A}$ and n -ary partially Hom-associativity of $\mathcal{A}'$, we obtain for $x_1, \dots, x_{2n-1} \in \mathcal{A}$ and $x'_1, \dots, x'_{2n-1} \in \mathcal{A}'$.

$$\begin{aligned}
& \sum_{i=0}^{n-1} m \otimes m'(\alpha_1 \otimes \alpha'_1(x_1 \otimes x'_1), \dots, \alpha_i \otimes \alpha'_i(x_i \otimes x'_i), m \otimes m'(x_{i+1} \otimes x'_{i+1}, \dots, x_{i+n} \otimes x'_{i+n}), \\
& \quad \alpha_{i+1} \otimes \alpha'_{i+1}(x_{i+1+n} \otimes x'_{i+1+n}), \dots, \alpha_{n-1} \otimes \alpha'_{n-1}(x_{2n-1} \otimes x'_{2n-1})) \\
&= \sum_{i=0}^{n-1} m(\alpha_1(x_1), \dots, \alpha_i(x_i), m(x_{i+1} \dots, x_{i+n}), \alpha_{i+1}(x_{i+1+n}), \dots, \alpha_{n-1}(x_{2n-1})) \otimes \\
& \quad m'(\alpha'_1(x'_1), \dots, \alpha'_i(x'_i), m'(x'_{i+1}, \dots, x'_{i+n}), \alpha'_{i+1}(x'_{i+1+n}), \dots, \alpha'_{n-1}(x'_{2n-1})) \\
&= m(\alpha_1(x_1), \dots, \alpha_j(x_j), m(x_{j+1} \dots, x_{j+n}), \alpha_{j+1}(x_{j+1+n}), \dots, \alpha_{n-1}(x_{2n-1})) \otimes \\
& \quad \sum_{i=0}^{n-1} m'(\alpha'_1(x'_1), \dots, \alpha'_i(x'_i), m'(x'_{i+1}, \dots, x'_{i+n}), \alpha'_{i+1}(x'_{i+1+n}), \dots, \alpha'_{n-1}(x'_{2n-1})) = 0
\end{aligned}$$

□

Proposition 3.3. *Let $\mathcal{A}$ be a symmetric n -ary totally Hom-associative algebra and $\mathcal{B}$ be an n -ary Lie algebra, then $\mathcal{A} \otimes \mathcal{B}$ is an n -ary Lie algebra.*

Proof. The proof is similar to the proof of the previous proposition. □

REFERENCES

- [1] Abramov V., Le Roy B. and Kerner R., *Hypersymmetry : a Z3-graded generalization of supersymmetry*, Journ. in Math. Phys., **38** (3), 1650-1669, (1997).
- [2] Aizawa, N., Sato, H., *q-deformation of the Virasoro algebra with central extension*, Physics Letters B, Phys. Lett. B **256**, no. 1, 185–190 (1991). Hiroshima University preprint preprint HUPD-9012 (1990).
- [3] Ataguema H. and Makhlouf A., *Deformations of Ternary Algebras*, J. Gen. Lie Theory and App. **1** (2007), 41-55.
- [4] Ataguema H. and Makhlouf A., *Notes on Cohomologies of Ternary Algebras of Associative Type*, arXiv:0812.0707v2 [math.RA] (2008).
- [5] Bagger J. and Lambert N., *Gauge Symmetry and Supersymmetry of Multiple M2-Branes*, arXiv:0711.0955, (2007).
- [6] Bazunova N., Borowiec A., and Kerner R., *Universal differential calculus on ternary algebras*, Lett. Math. Phys. **67**, 195-206 (2004).
- [7] Bremmer M.R. and Peresi L.A., *Ternary analogues of Lie and Malcev algebras*, Linear Algebra and its Applications **414**, 1–18, (2006).
- [8] Borowiec A., Dudek W. and Duplij S., *Basic concepts of ternary Hopf algebras*, Journal of Kharkov National University, ser. Nuclei, Particles and Fields, V. **529** - N 3 (15), 21–29, (2001).
- [9] Carlsson R., *N-ary algebras*, Nagoya Math. J. **78**, 45-56, (1980).
- [10] Carlsson R., *Cohomology of associative triple systems*, Proc. Amer. Math. Soc. **60**, 1-7, (1976).
- [11] Cassas J.M., Loday J.-L. and Pirashvili, *Leibniz n-algebras*, Forum Math. **14**, 189–207, (2002).
- [12] Chaichian M., Ellinas D. and Z. Popowicz, *Quantum conformal algebra with central extension*, Phys. Lett. B **248**, no. 1-2, 95–99 (1990).
- [13] Chaichian M., Isaev A. P., Lukierski J., Popowicz Z. and Prešnajder P., *q-deformations of Virasoro algebra and conformal dimensions*, Phys. Lett. B **262** (1), 32–38 (1991).
- [14] Chaichian M., Kulish P. and Lukierski J., *q-deformed Jacobi identity, q-oscillators and q-deformed infinite-dimensional algebras*, Phys. Lett. B **237**, no. 3-4, 401–406 (1990).
- [15] Chaichian M., Popowicz Z. and Prešnajder P., *q-Virasoro algebra and its relation to the q-deformed KdV system*, Phys. Lett. B **249**, no. 1, 63–65 (1990).
- [16] Curtright, T. L., Zachos, C. K., *Deforming maps for quantum algebras*, Phys. Lett. B **243**, no. 3, 237–244 (1990).
- [17] Damaskinsky E. V. and Kulish P. P., *Deformed oscillators and their applications*, (Russian), Zap. Nauch. Semin. LOMI **189**, 37–74 (1991); Engl. transl. in J. Soviet Math. **62** (5), 2963–2986 (1992).
- [18] Daskaloyannis C., *Generalized deformed Virasoro algebras*, Modern Phys. Lett. A **7** no. 9, 809–816 (1992).
- [19] De Azcarraga J. A., Izquierdo J. M., and Perez J. C. Bueno, *On the higher-order generalizations of Poisson structures*, J. Phys. A: Math. Gen. **30**, 607–616 (1997).
- [20] De Azcarraga J. A., Perelomov A. M., and Perez Bueno J. C., *The Schouten-Nijenhuis bracket, cohomology and generalized Poisson structures*, J. Phys. A: Math. Gen. **29**, 7993–8009 (1996).
- [21] De Azcarraga J. A. and Perez Bueno J. C., *Multibracket simple Lie algebras*, In “Physical Applications and Mathematical Aspects of Geometry, Groups and Algebra”, Vol I, pp. 103–107. World Sci., 1997.

- [22] Dito G., Flato M., Sternheimer D. and Takhtajan L., *Deformation Quantization and Nambu Mechanics*, Commun. Math. Phys., **183**, 1–22. (1997)
- [23] Dito G., Flato M., Sternheimer D., *Nambu mechanics, n -ary operations and their quantization*, in: Deformation Theory and Symplectic Geometry, Mathematical Physics Studies, vol. **20**, Kluwer Acad. Publ., 43–66, (1997).
- [24] Filippov V. T., *n -ary Lie algebras*, Sibirskii Math. J. **24**, 126–140 (1985).
- [25] Gautheron P., *Some Remarks Concerning Nambu Mechanics*, Letters in Mathematical Physics **37**: 103–116, (1996).
- [26] Gautheron P., *Simple facts concerning Nambu algebras*, Com. in Mathematical Physics **195** (2), 417–434 (1998).
- [27] Gnedbaye A. V., *Les algèbres K -aires et leurs opérades*, C. R. Acad. Sci. Paris. série I, **321**, 147–152 (1995).
- [28] Gnedbaye A. V., *Opérades, algèbres de Leibniz et triples de Jordan*, Mémoire de synthèse pour l’habilitation à diriger des recherches, Preprint IRMA Strasbourg (2003).
- [29] Goze M. and Remm E., *Lie-admissible coalgebras*, J. Gen. Lie Theory Appl. **1** (1), 19–28 (2007).
- [30] Goze N. and Remm E., *On n -ary algebras given by Gerstenhaber’s products*, arXiv:0803.0553v2 [math.RA], (2008).
- [31] Goze N. and Rausch de Traubenberg M., *Hopf algebras for ternary algebras and groups*. arXiv:0809.4212 (2008).
- [32] Grabowski J. and Marmo G., *Remarks on Nambu-Poisson and Nambu-Jacobi brackets*, J. Phys. A: Mah. Gen. **32**, 4239–4247 (1999).
- [33] Grabowski J., *Abstract Jacobi and Poisson structures*, J. Geom. Phys. **9**, 45–73 (1992).
- [34] Hanlon Ph. and Wachs M., *On Lie k -Algebras*, Adv. Math. **113**, 206–236 (1995).
- [35] Harris B., *Cohomology of Lie triple systems and Lie algebras with involution*, Trans. Amer. Math. Soc **98**, 148–162 (1961).
- [36] Hartwig J. T., Larsson D. and Silvestrov S. D., *Deformations of Lie algebras using σ -derivations*, J. of Algebra **295**, 314–361 (2006).
- [37] Hellström L. and Silvestrov S. D., *Commuting elements in q -deformed Heisenberg algebras*, World Scientific, 2000.
- [38] Hestenes M. R., *On ternary algebras*, Scripta Math. **29**, no. 3-4, 253–272 (1973).
- [39] Hoffbeck E., *A Poincar-Birkhoff-Witt criterion for Koszul operads*. arXiv:0709.2286v3 [math.AT] (2007).
- [40] Hu N., *q -Witt algebras, q -Lie algebras, q -holomorph structure and representations*, Algebra Colloq. **6**, no. 1, 51–70 (1999).
- [41] Ibanez R., de Leon M. J., Marrero C. and Martin de Diego D., *Dynamics of generalized Poisson and Nambu-Poisson brackets*, J. Math. Phys. **38** (5), 2332–2344 (1997).
- [42] Jacobson N., *Lie and Jordan triple systems*, Amer. J. Math. **71**, 149–170 (1949).
- [43] Kapranov M., Gelfand M. and Zelevinskii A., *Discriminants, Resultants and Multidimensional Determinants*, Berlin Birkhauser (1994).
- [44] Kassel C., *Cyclic homology of differential operators, the Virasoro algebra and a q -analogue*, Commun. Math. Phys. **146**, 343–351 (1992).
- [45] Katsylo P. and Mikhailov D., *Ternary quartics and 3-dimentional commutative algebras*, arXiv:alg-geom/9408900 (1994).
- [46] Kerner R., *Ternary algebraic structures and their applications in physics*, in the “Proc. BTLP 23rd International Colloquium on Group Theoretical Methods in Physics”, ArXiv math-ph/0011023, (2000).
- [47] Kerner R., *Z_3 -graded algebras and non-commutative gauge theories*, dans le livre “Spinors, Twistors, Clifford Algebras and Quantum Deformations”, Eds. Z. Oziewicz, B. Jancewicz, A. Borowiec, pp. 349-357, Kluwer Academic Publishers (1993).
- [48] Kerner R., *Z_3 -grading and ternary algebraic structures*, dans les Proceedings du Workshop “New Symmetries and Differential Geometry”, Clausthal 1993, V. Dobrev, M.D. Doebner and S. Ushveridze eds., pp. 375-394, World Scientific (1994).
- [49] Kerner R., *The cubic chessboard : Geometry and physics*, Classical Quantum Gravity **14**, A203-A225 (1997).
- [50] Kerner R., *Z_3 -graded ternary algebras, new gauge theories and quarks*, In Proceedings du Workshop “Topics in Quantum Field Theory”, Maynooth 1995, ed. T. Tchrakian, World Scientific, 113–126 (1995).
- [51] Kerner R. and Vainerman L., *On special classes of n -algebras*, Journ. in Math. Phys., **37** (5), 2553–2565, (1996).
- [52] Kerner R., *Generalized Cohomologies and Differentials of Higher Order*, Proceedings de la conference DGMTP de Tianjin (Chine),. Eds. Ge, Wang, World Scientific (2006).
- [53] Kurosh A. G., *Multioperator rings and algebras*, Russ Math Surv, **24** (1), 1–13 (1969).
- [54] Larsson D. and Silvestrov S. D., *Quasi- Hom-Lie algebras, Central Extensions and 2-cocycle-like identities*, J. of Algebra **288**, 321–344 (2005).
- [55] Larsson D. and Silvestrov S. D., *Quasi-Lie algebras*, in “Noncommutative Geometry and Representation Theory in Mathematical Physics”, Contemp. Math., **391**, Amer. Math. Soc., Providence, RI , 241–248 (2005).
- [56] Larsson D. and Silvestrov S. D., *Quasi-deformations of $sl_2(\mathbb{F})$ using twisted derivations*, Comm. in Algebra **35** , 4303 – 4318 (2007).
- [57] Lister W. G., *Ternary rings*, Trans. Amer. Math. Soc. **154**, 37–55 (1971).
- [58] Liu Ke Qin, *Characterizations of the quantum Witt algebra*, Lett. Math. Phys. **24** , no. 4, 257–265 (1992).
- [59] Loos O., *Assoziative tripelsysteme*, Manuscripta Math. **7**, 103–112 (1972).
- [60] Loos O., *Symmetric spaces*, vol. **1**, 103–112, W. A. Benjamin, New York (1969).
- [61] Makhlouf A. and Silvestrov S. D., *Hom-algebra structures*, J. Gen. Lie Theory Appl. **2** (2) , 51–64 (2008).

- [62] Makhlouf A. and Silvestrov S. D., *Hom-Lie admissible Hom-coalgebras and Hom-Hopf algebras*, Published as Chapter 17, pp 189-206, S. Silvestrov, E. Paal, V. Abramov, A. Stolin, (Eds.), Generalized Lie theory in Mathematics, Physics and Beyond, Springer-Verlag, Berlin, Heidelberg, (2008).
- [63] Makhlouf A. and Silvestrov S. D., *Notes on Formal deformations of Hom-Associative and Hom-Lie algebras*, To appear in Forum Mathematicum. Preprints in Mathematical Sciences, Lund University, Centre for Mathematical Sciences, Centrum Scientiarum Mathematicarum, (2007:31) LUTFMA-5095-2007, 2007; arXiv:0712.3130v1 [math.RA] (2007).
- [64] Makhlouf A. and Silvestrov S. D., *Hom-Algebras and Hom-Colgebras*, Preprints in Mathematical Sciences, Lund University, Centre for Mathematical Sciences, Centrum Scientiarum Mathematicarum, (2008) ; arXiv:0811.0400 [math.RA] (2007).
- [65] Michor P. W. and Vaisman I. , *A note on n -ary Poisson brackets*, Proceedings of the 19th Winter School "Geometry and Physics" (Srní, 1999). Rend. Circ. Mat. Palermo (2) Suppl. No. **63** , 165–172 (2000).
- [66] Michor P. W. and Vinogradov A. M., *n -ary Lie and associative algebras*, Rend. Sem. Mat. Univ. Pol. Torino. **54**, 373-392 (1996).
- [67] Myung H. C., *A characterization of the Jacobson radical in ternary algebras*, Proc. Amer. Math. Soc. **38**, 228–234 (1973).
- [68] Nambu Y., *Generalized Hamiltonian mechanics*, Phys. Rev. **D7**, 2405–2412 (1973).
- [69] Okubo S. *Triple products and Yang-Baxter equation (I): Octonionic and quaternionic triple systems*, J. Math.Phys. **34**, 3273–3291 (1993).
- [70] Rausch de Trautenberg M. and Slupinski M. J., *Finite-dimensional Lie algebras of order F* , J. Math. Phys. **43**, 5145–5160 (2002).
- [71] Rausch de Trautenberg., *Ternary algebras and groups*. J. Phys.: Conf. Ser. 128, (2008).
- [72] Rausch de Trautenberg., *Some Results on Cubic and Higher Order Extensions of the Poincar Algebra*, arXiv:0811.1465v1 [hep-th], (2008).
- [73] Remm. E., *On the NonKoszulity of $(2p+1)$ -ary partially associative Operads*, arXiv:0812.2687v1 [math.AT] (2008).
- [74] Santana A. E. and Muradian R., *Hopf Structure in Nambu-Lie n -Algebras*, Theoretical and Mathematical Physics, **114** (1), (1998).
- [75] Sokolov N. P., *Introduction to the theory of Multidimensional Matrices*, Kiev Naukova Dumaka(1972).
- [76] Takhtajan L., *On foundation of the generalized Nambu mechanics*, Comm. Math. Phys. **160** , 295-315 (1994).
- [77] Takhtajan L., *A higher order analog of Chevalley-Eilenberg complex and deformation theory of n -algebras*, St. Petersburg Math. J. **6**, 429–438 (1995).
- [78] Yamaguti K., *On the cohomologie space of Lie triple systems*, Kumamoto J. Sci. Ser. A **5**, 44–52(1960).
- [79] Yau D., *Enveloping algebra of Hom-Lie algebras*, J. Gen. Lie Theory Appl. **2** (2), 95–108 (2008).
- [80] Yau D., *Hom-algebras as deformations and homology*, arXiv:0712.3515v1 [math.RA] (2007).
- [81] Yau D., *Hom-bialgebras and comodule algebras*, arXiv:0810.4866v1 [math.RA] (2008).

ABDENACER MAKHLOUF AND HAMMIMI ATAGUEMA, UNIVERSITÉ DE HAUTE ALSACE, LABORATOIRE DE MATHÉMATIQUES, INFORMATIQUE ET APPLICATIONS, 4, RUE DES FRÈRES LUMIÈRE F-68093 MULHOUSE, FRANCE
E-mail address: Abdenacer.Makhlouf@uha.fr

SERGEI SILVESTROV, CENTRE FOR MATHEMATICAL SCIENCES, LUND UNIVERSITY, BOX 118, SE-221 00 LUND, SWEDEN
E-mail address: sergei.silvestrov@math.lth.se